

Chiral-dispersive calculations of $\pi\pi$ scattering confront experiment*

J.R.Peláez ^a and F. J. Ynduráin^b

^a Dept. de Física Teórica II, Univ. Complutense de Madrid. 28040 Madrid. Spain

^b Dept. de Física Teórica, C-XI Univ. Autónoma de Madrid, Canto Blanco, E-28049, Madrid, Spain.

In a series of papers we have applied several sum rules and forward dispersion relations, to $\pi\pi$ scattering. We have found that some widely used data sets fail to satisfy these constraints, and we have provided an amplitude that describes data consistently with the dispersive tests. Furthermore, we noted that the input and precision claimed in a Roy equation analysis by Colangelo, Gasser and Leutwyler (CGL), lead to several mismatches with some sum rules. Subsequently, Caprini, Colangelo, Gasser and Leutwyler claimed that our Regge parametrization was incorrect. We collect here the answers to their various claims, and try to clarify the points of agreement and disagreement, showing *experimental* evidence that substantiates our results, and that, in addition, their representation fails to satisfy all three forward dispersion relations up to $\sqrt{s} \leq 800$ MeV by several standard deviations.

Introduction

Some time ago, Ananthanarayan, Colangelo, Gasser and Leutwyler [1] (ACGL) and Colangelo, Gasser and Leutwyler [2] (CGL) have used experimental information, unitarity, analyticity (in the form of the Roy equations) and, in CGL, chiral perturbation theory, to construct the $\pi\pi$ scattering amplitude. In CGL, an outstanding precision was claimed, at the percent level, for scattering lengths and effective ranges of S, P, D and F waves. In addition, CGL provided parametrizations for the S, P phase shifts up to $s^{1/2} = 0.8$ GeV.

In a series of works, referred to as PY1,[3] PY-Regge,[4] PY-Sardinia[5] and PY-FDR[6] (collectively, PY) we have contested the *input* of both ACGL and CGL for the following reasons:

- i. The high energy ($s^{1/2} > 1.4$ GeV), in particular the Regge parametrization, is incompatible with data and violates factorization.
- ii. A relevant source of uncertainty, the error on the phase $\delta_0^{(0)}$ at 800 MeV, a crucial input for the matching between low and high energy representations, is largely underestimated. The

central value of either $\delta_0^{(0)}$ or $\delta_0^{(2)}$ is inconsistent with forward dispersion relations.

- iii. The D2 wave disagrees, both at low and high energy, with what is found from experiment.

This shed doubts on the precision claimed in CGL (final uncertainties in ACGL are larger). In fact, using experimental input, we showed that

- iv. Some low energy parameters given in CGL, mainly those for D waves, do not satisfy the Froissart Gribov sum rules. The P wave effective range parameter, b_1 , deviates by several standard deviations from what one finds from the pion form factor [7] or from a sum rule.
- v. The CGL final amplitudes fail to verify forward dispersion relations up to 800 MeV, again by several standard deviations.

In the time between PY1 and PY-Regge, Caprini, Colangelo, Gasser and Leutwyler[8] (CCGL) claimed to counter the criticism in PY1. These claims were answered in PY-Regge and PY-FDR (and PY-Sardinia and [9]). However, in this note we provide a more formal response collecting all arguments together and also including comments on a recent article of Colangelo [10].

Let us first enumerate the main points in which CCGL claim to answer the criticism of PY1:

*Joint contribution of the two talks presented by the authors in QCD05, Montpellier, France, July 2005.

1. The asymptotic behaviour used in PY1. CCGL still consider it inferior to that used in ACGL and CGL in that it is not in “equilibrium” with low energy, and that our asymptotics “violates crossing rather strongly”.
2. Based only on a selection of data, CCGL conclude that CGL describes the S0 data, whereas our “tentative solution” does not.
3. CCGL claim that, even with PY1 Regge formulas, the CGL results are essentially unchanged.
4. CCGL consider the *Olsson sum rule* and agree that it is not satisfied by CGL, if using the Regge asymptotics of PY1. From this, CCGL conclude that “the asymptotics used in PY1 is inconsistent with the theoretical predictions for S-wave scattering lengths”.
5. They keep their b_1 prediction claiming that the Froissart Gribov sum rule is subject to large uncertainties due to the Regge representation.

1. Regge asymptotics and the D2 wave

The ACGL and CGL Regge violates factorization, a well-known property of Regge theory, for all trajectories, but, in particular, for the Pomeron exchange by a factor larger than two.

The Regge parameters in ACGL, CGL are obtained by “balancing” the high and low energy contributions in a number of crossing sum rules. CCGL conclude that the Regge representation in PY1 is *incompatible with crossing symmetry* because it does not satisfy crossing sum rules, that are better satisfied by their Regge representation (which they concede that could be improved).

However, although ignored then by ACGL, CGL, CCGL and ourselves (we remedied this in PY-Regge), there are many data on $\pi^0\pi^-$ and, particularly, on $\pi^-\pi^-$ and $\pi^+\pi^-$ total cross sections at high energy [11]. This is shown in Fig. 1 where it is clearly seen that the Regge expressions used by ACGL, CGL and CCGL are systematically below the data whereas our parametrization falls on top of them. In fact, in PY-Regge, we used the $\pi\pi$ experimental data, together with the very precise πN , KN and NN cross sections data to get an accurate verification of factorization




Figure 1 $\pi\pi$ total cross section data [11]. The stars (PY) are reconstructed in PY-Regge from experimental phase shift analysis. Continuous lines (PY): Regge parameters as in PY-Regge, in agreement, within errors with PY1. Dashed lines: ACGL Regge representation. The grey bands cover the respective uncertainties. The dotted line is the reconstructed $\pi^+\pi^-$ cross section from the Cern–Munich analysis; (Fig. 7 in [14])

and a precise Regge parametrization. In the Appendix B of PY-FDR, by means of the Froissart Gribov representation, we refined the ρ residue and slope obtaining a value compatible with our previous determinations.

Although the data supports our Regge parametrization, one might still wonder about the sum rules. However [9] these sum rules are (A) fairly well satisfied by the PY1 representation, *when errors are taken into account*, which are notoriously absent in Eqs. (11) and the next equation in CCGL; and, (B) At low energies the S-wave contribution cancels, and, in some cases, also the P wave is absent. Thus, these sum rules are dominated by the D waves. But, for the D2 wave, ACGL and CGL borrow the old fit from the book of Martin, Morgan and Shaw [12]

$$\delta_2^{(2)}(s) = -0.003(s/4M_\pi^2)(1 - 4M_\pi^2/s)^{5/2}. \quad (1)$$

Although it was obtained only from data in the $0.625\text{GeV} \leq s^{1/2} \leq 1.375\text{GeV}$ region, ACGL and CGL use it from threshold up to 2 GeV.

In fact, (1) gives a negative scattering length, whereas it is known that $a_2^{(2)}$ is *positive*, and it does not fit well the data below 0.7 GeV, as shown

in Fig. 2. Above 1 GeV, the modulus of (1) grows like s , whereas Regge theory requires all waves to tend to a multiple of π ; i.e. D2 should go to zero. In addition, we see in Fig. 2 that Eq.(1) does not fit well the data between 1.4 and 2 GeV. Finally, above 1.5 GeV, where the $\pi^-\pi^-\rightarrow\rho^-\rho^-$ channel opens, the D2 wave should be highly inelastic: but ACGL take it elastic up to 2 GeV. The fact that the Regge representation of ACGL, CGL, fits their sum rules is more *negative* than positive support. For further detail, we refer to [9], PY1 and PY-FDR where we show that crossing sum rules are perfectly satisfied when considering the correct Reggeistics *and* the correct D waves.




Figure 2. $I = 2$, D -wave phase shift and data [13]. Continuous line: (PY-FDR) fit to data. Dashed line: (PY-FDR) improved parameters using dispersion relations. Dotted line: Martin, Morgan and Shaw fit which ACGL, CGL and CCGL use from threshold to $s^{1/2} = 2$ GeV.

2. CCGL claim the data are described by CGL but not by our “tentative solution”

In PY1 we presented a “tentative solution”, merely a fit to low energy $\pi\pi$ data, in order to compare with the CGL low energy partial waves. CCGL conclude that this tentative solution does not fit the experimental data, at least not as well as the phase shifts in CGL do. This is very surprising since, our “tentative solution” was obtained by just fitting data. We will discuss here the S0 wave, and particularly the value of its phase shift at 800 MeV, a key value in their input, and then the S2 wave.

2.1. The S0 wave below 800 MeV

By looking at Fig. 1 in CCGL, reproduced here in Fig. 3a only with the CGL and “tentative solution” of PY1, it may seem that data fall on the CGL results, and are incompatible with the PY1 “tentative solution”. However, *that only happens because CCGL have not plotted all data*. Fig. 1 of CCGL is certainly unfair with our tentative solution, which is a fit to an average of *published* data. Indeed, in Fig. 3b here, we include the solutions of different experimental analyses [14,15,16,17,18,20] which ACGL, CGL quote in their references, but CCGL do not show in their figure (we have also included in Fig. 3b the recent data in [19]). It is not clear why ACGL, CGL and CCGL only consider a subset of all published data.




Figure 3. S0 phase shifts. a) The shaded area is the CGL error band. The PY1 tentative solution lies between the dashed lines. Only the data included in Fig. 1 of CCGL are shown. b) PY1 Tentative solution and improved solution of PY-FDR. The dashed line is the CGL phase. We show data from [14,15,16,17,18,19,20]. (see PY-FDR for

details)

2.2. The matching phase $\delta_0^{(0)}(0.8 \text{ GeV})$

One of the most important input parameters [10] in ACGL, CGL is the S0 phase shift at the matching point between low and high energy, $s_m \equiv (0.8 \text{ GeV})^2$. The data is affected by large systematic errors, but ACGL consider the difference between S0 and P phase shifts $\delta_1(s_m) - \delta_0^{(0)}(s_m)$ hoping some uncertainties may cancel. In particular, by interpolation, they quote

$$\begin{aligned} 24.8 \pm 3.8^\circ & \quad [\text{Estabrooks and Martin, } s\text{-channel}] \\ 30.3 \pm 3.4^\circ & \quad [\text{Estabrooks and Martin, } t\text{-channel}] \\ 23.4 \pm 4.0^\circ & \quad [\text{Hyams et al.}] \end{aligned} \quad (2)$$

and average them to obtain $26.6^\circ \pm 3.7^\circ$.

However, these three numbers stem from different analyses of the *same experiment*, CERN-Munich [14,18], so that their spread measures the *systematic* error. Consequently, one should have enlarged the error to cover *all* central values:

$$\delta_1(s_m) - \delta_0^{(0)}(s_m) = 26.3^\circ \pm 3.8 (\text{sta.}) \pm 4.0 (\text{sys.}).$$

Furthermore, since Estabrooks and Martin,[18], in their section 4, state that the input uncertainties, particularly in the D-wave, lead to “*systematic changes in δ_S^0 of the order of 10°* ”, that is, the ± 4 systematic error is likely still optimistic.

Then ACGL average again their number with 26.5 ± 4.2 , interpolated from Protopopescu et al. [20] (Table VI), to obtain a surprisingly small error: $\delta_1(s_m) - \delta_0^{(0)}(s_m) \stackrel{\text{ACGL}}{=} 26.6^\circ \pm 2.8^\circ$. However, in Protopopescu et al. [20], we read that “*the given errors...reflect only statistical error...and...should be considered only as an indication of the minimum error in our computed values*”. Still, CGL add back $\delta_1(s_m) = 108.9 \pm 2^\circ$, and come up with,

$$\delta_0^{(0)}(0.8 \text{ GeV}) = 82.3 \pm 3.4^\circ, \quad (3)$$

which is also used in CGL and defended in [10].

To our view, the statistical and systematic errors should be added linearly, in order to be conservative, given the caveats of the original authors. *But even this is most likely still optimistic*, since there are more data, even more spreaded. In fact, the value from Hyams et al., [14] quoted

in (2), is only one of *five solutions published by the same experiment*, cf. Grayer et al.[15]. The datum of Protopopescu et al., is again one of the several analysis in [20], which differ among themselves again by about 10° – not by chance in agreement with the statements quoted above by Estabrooks and Martin, [18].




Figure 4. a) Values of $\delta_0^{(0)}(0.8 \text{ GeV})$, as given by Colangelo in Fig. 3 in ref 10. We include only data with error bars. The shaded area covers the error band in ACGL, Eq.(3) here. Systematic errors estimated [17] to be $\sim 10^\circ$ were not included. b) Published data [14,15,16,17,18,19,20] on $\delta_0^{(0)}(0.8 \text{ GeV})$. The band is like in 5.a.

Authors and references	$\delta_0^{(0)}(0.8 \text{ GeV})$
Kaminski et al. [19]	$98.3 \pm 5.3^\circ$
Grayer et al.,A [15]	$75.4 \pm 3.0^\circ$
Grayer et al.,B [15]	$81.7 \pm 3.9^\circ$
Grayer et al.,C [15]	$88.9 \pm 4.0^\circ$
Grayer et al.,E [15]	$69.5 \pm 3.8^\circ$
EM,s-channel [18]	$90.4 \pm 3.6^\circ$
EM,t-channel [18]	$85.7 \pm 2.9^\circ$
Protopopescu et al.[20] (VI)	$81.6 \pm 4.0^\circ$
Protopopescu et al. [20](VIII)	$73.4 \pm 4.0^\circ$

Table 1. $\delta_0^{(0)}$ values *interpolated* to 0.8 GeV from different experimental analysis. ACGL, CGL and CCGL, take as experimental input , $\delta_0^{(0)}(0.8, \text{GeV}) = 82.3 \pm 3.4^\circ$.

The ACGL, CGL matching input is summarized by Colangelo in [10], where in his Fig. 3 the ACGL uncertainty band is shown, but compared *only* with the data used in CGL and ACGL, and *only with statistical errors*. This can be seen in our Fig.4a above, where we plot the ACGL band, Eq.(3) here, with the data included in Fig. 3 of [10]. Here we do not include the data that does

not have error bars, since, they are not even used by ACGL to estimate their uncertainty. In contrast, in Fig. 4b we repeat Fig. 4a, but we add other *published* data sets. We list the actual values and references in Table 1: The small error band of ACGL does not represent the experimental uncertainty. Even less so when considering that many data come from different analyses of the same experiment (like CERN-Munich and Estabrooks and Martin), and their central value spread is not of statistical nature.

Finally, we want to remark that without imposing the ACGL, CGL surprisingly small uncertainty on the matching phase, a simultaneous fit [21] to Roy equations and data yields $\delta_0^{(0)} = 92.6^\circ$ whereas a fit only to data gives $\delta_0^{(0)} = 87.2^\circ$, both outside the value imposed by CGL and CGL as their input.




Figure 5. The $I = 2$, S -wave. Data comes from [4,13]. Continuous line: PY-FDR fit. Dotted line: PY1 fit. Data from Durusoy et al. and from Solution B of Hoogland et al. were not included in the fits (see PY-FDR for details). The dashed line, below our fit, is the S2 phase of CGL [2]. The gray bands cover the respective uncertainties.

2.3. The S2 wave

The S2 phase shift data [13] can be fitted with a simple effective range expansion, like the “tentative solution” of PY1, where we neglected the inelasticity below 1450 MeV, or, with greater precision, in PY-FDR, including the inelasticity above ~ 1 GeV. In Fig. 5 the PY fit is shown to describe data better than the CGL phase shift: not surprisingly since it was *not* obtained from theory

as the CGL solution. In fact, the χ^2 of the latter gets twice as large as that of the PY1 and PY-FDR fits, as $\sqrt{s}$ tends to 800 MeV. In addition, we show below that forward dispersion relations are in conflict with a curve as low as that of CGL.

3. Regge formulas and threshold results

CCGL repeated the Roy equation analysis of CGL using the Regge PY1 formulas, and claimed that their results, except for the S2 wave, do not vary appreciably. Let us first note that Roy equations use Regge expressions beyond their applicability region $|t| \ll s$. Second, that the effect of Regge formulas is strongly constrained since CGL are forcing their solutions to match the $\delta(s_m)$ phase shift with an extremely small input uncertainty that, as we have seen in the previous section, basically neglects all systematic uncertainties.

However, low energy parameters can be calculated [3] with sum rules that involve only small values of $|t|$: First, $a(+0) \equiv a_2(\pi^0\pi^0 \rightarrow \pi^+\pi^-) = 2[a_2^{(0)} - a_2^{(2)}]/3$, and $a(00) \equiv a_2(\pi^0\pi^0 \rightarrow \pi^0\pi^0) = 2[a_2^{(0)} + 2a_2^{(2)}]/3$ can be calculated with the Froissart Gribov (FG) representation, that needs only information at $t = 4M_\pi^2 \simeq 0.08 \text{ GeV}^2$. Second, $2a_0^{(0)} - 5a_0^{(2)}$, can be evaluated with the Olsson sum rule, that just needs $t = 0$. We find that

i) Using inside the sum rule integrals the low energy parameterizations of CGL up to 800 MeV, and experiment between 800 and 1.42 GeV, together with the Regge parameters of PY1, we obtain $a(0+) = 10.94 \pm 0.13$. Thus, the CGL result, $a(0+) = 10.53 \pm 0.10$, obtained with a Wanders sum rule, using their Regge and D-wave parameterizations instead, presents a 2.5σ mismatch. For the difference between the CGL calculation using Wanders sum rules minus the FG representation, which cancels correlations, the mismatch is of more than 4 standard deviations. We agree with CCGL that this difference does not involve the S and P waves, and the mismatch is only due to the Regge and $l \geq 2$ wave input, different for CGL and PY. However, as we have just seen, it certainly affects the $a(0+)$ total value by about 2.5σ . The situation for $a(00)$ is very similar [3].

ii) From Eq. (11.2) in CGL, we find, in units of the pion mass,

$$2a_0^{(0)} - 5a_0^{(2)} = 0.663 \pm 0.006 \quad \text{“CGL, direct”} \quad (4)$$

Alternatively, we can use the Olsson sum rule:

$$2a_0^{(0)} - 5a_0^{(2)} = 3M_\pi \int_{4M_\pi^2}^{\infty} ds \frac{\text{Im } F^{(I_t=1)}(s, 0)}{s(s - 4M_\pi^2)}. \quad (5)$$

The total $I_t = 1$ Regge contribution comes out the same either with the PY1 parametrization, or with the improved PY-Regge and PY-FDR Regge parameters, i.e. replacing the contribution of the ρ' trajectory and of the $\rho(1450)$ resonance for a slight increase in the ρ residue to $\beta_\rho(0) = 1.02 \pm 0.11$. At low energy we use the S, P waves of CGL. We obtain 0.635 ± 0.014 , in pion mass units. There is a clear mismatch between the “direct” result, Eq.(4), and the dispersive evaluations of $2a_0^{(0)} - 5a_0^{(2)}$, which can be calculated with large precision. Concerning the S and P wave scattering lengths individually, the PY and CGL results are in agreement, mostly due to the larger PY error bars (thus, our results do not question the large condensate scenario of ChPT).

In summary, some quantities may not change if using different Regge asymptotics, but certainly others do.

4. CGL solutions and forward dispersion relations

In PY-FDR we studied a forward dispersion relation for the $I_t = 1$, which at threshold reduces to the Olsson sum rule already evaluated in PY1. In addition, we studied subtracted forward dispersion relations for $\pi^0\pi^+$ and $\pi^0\pi^0$ scattering. Indeed, these relations imply the vanishing of:

$$\begin{aligned} \Delta_1 &\equiv \text{Re } F^{(I_t=1)}(s, 0) \\ &- \frac{2s - 4M_\pi^2}{\pi} \text{P.P.} \int_{4M_\pi^2}^{\infty} ds' \frac{\text{Im } F^{(I_t=1)}(s', 0)}{(s' - s)(s' + s - 4M_\pi^2)}, \\ \Delta_{00} &\equiv \text{Re } F_{00}(s) - F_{00}(4M_\pi^2) - \frac{s(s - 4M_\pi^2)}{\pi} \times \\ &\text{P.P.} \int_{4M_\pi^2}^{\infty} ds' \frac{(2s' - 4M_\pi^2) \text{Im } F_{00}(s')}{s'(s' - s)(s' - 4M_\pi^2)(s' + s - 4M_\pi^2)}, \end{aligned}$$

$$\begin{aligned} \Delta_{0+} &\equiv \text{Re } F_{0+}(s) - F_{0+}(4M_\pi^2) - \frac{s(s - 4M_\pi^2)}{\pi} \times \\ &\text{P.P.} \int_{4M_\pi^2}^{\infty} ds' \frac{(2s' - 4M_\pi^2) \text{Im } F_{0+}(s')}{s'(s' - s)(s' - 4M_\pi^2)(s' + s - 4M_\pi^2)} \end{aligned}$$

We plot these differences in Fig. 6, using the CGL parameters for the S0, S2 and P waves below 800 MeV where the mismatch is clearly seen. In contrast, as shown in PY-FDR, forward dispersion relations are fairly well satisfied below 800 MeV by simple fits to data on S0, S2 and P waves, and remarkably well for our improved PY-FDR solutions.




Figure 6 Continuous lines: for CGL S and P phase shifts, the differences δ_i between the real parts calculated directly or from the dispersive formulas. Consistency occurs when curves fall within the shaded areas [5].

Finally, one may wonder why there is a mismatch for CGL, since their S0 wave follows the Solution B of Grayer et al.[15], whose improved fit is fairly compatible with dispersion relations (with the correct Regge behaviour and D2 wave,

see PY-FDR). However one should note that the Solution B fit, constrained to satisfy dispersion relations, leads to an amplitude that violates the $\pi^0\pi^0$ dispersion relation at $s = 2M_\pi^2$ (Table 3 in PY-FDR). In addition, it requires S2 phase shifts that disagree even more with that in CGL than the S2 wave obtained only by fitting data, as shown in Fig. 7. Indeed, the S2 wave obtained by CCGL from Roy equations but using PY asymptotics is also displaced in the same direction.




Figure 7 The $I = 2$, S -wave phase shift. Data as in Fig. 4. Continuous line: CGL solution and error band. Dotted line: PY-FDR fit. Dashed line: PY-FDR fit constrained with dispersion relations when also improving the S0 wave fit from Solution B of Grayer et al.[15].

5. The value of b_1 .

In PY1 it was noted that the CGL value for the effective range parameter, $b_1 = (5.67 \pm 0.13) \times 10^{-3} M_\pi^{-5}$, was several standard deviations away from what one gets from the pion form factor [7] $b_1 = (4.73 \pm 0.23) \times 10^{-3} M_\pi^{-5}$. CCGL answered that such a result depended on the specific form of the parametrization used for the P wave phase. To clarify this matter, in PY-FDR we devised a fastly convergent sum rule that depends little on the high energy behaviour or the low-energy P wave phase shift, so it provides an independent determination of b_1 :

$$M_\pi b_1 = \frac{2}{3} \int_{4M_\pi^2}^{\infty} ds \left\{ \frac{1}{3} \left[\frac{1}{(s - 4M_\pi^2)^3} - \frac{1}{s^3} \right] \text{Im } F^{(I_t=0)}(s, 0) \right.$$

$$\left. + \frac{1}{2} \left[\frac{1}{(s - 4M_\pi^2)^3} + \frac{1}{s^3} \right] \text{Im } F^{(I_t=1)}(s, 0) - \frac{5}{6} \left[\frac{1}{(s - 4M_\pi^2)^3} - \frac{1}{s^3} \right] \text{Im } F^{(I_t=2)}(s, 0) \right\}.$$

The largest contribution comes from S0 and P waves at low energy, while all other pieces (in particular, the Regge contributions) are substantially smaller than 10^{-3} . We find

$$b_1 = (4.99 \pm 0.21) \times 10^{-3} M_\pi^{-5}. \quad (6)$$

In PY-FDR, using fits constrained with dispersion relations, we found $b_1 = (4.55 \pm 0.21) \times 10^{-3} M_\pi^{-5}$. In conclusion, all three experiment-based values for b_1 are fairly compatible among themselves, but several standard deviations away from the CGL value.

Another matter addressed to by Colangelo in ref. 10 is that of the *scalar* form factor of the pion. We will not discuss this here, but refer to the relevant literature [22].

6. Conclusions

In a series of works we have contested the input used by Ananthanarayan, Colangelo, Gasser and Leutwyler [1] (ACGL) and Colangelo, Gasser and Leutwyler [2] in their Roy equation analysis of $\pi\pi$ scattering. This challenged the remarkable precision claimed in CGL. Subsequently, Caprini, Colangelo, Gasser and Leutwyler [8,10] claimed to refute our arguments. Here, we collect the answers to their arguments, clarifying the points of agreement and disagreement, and showing *experimental* support for our results. In particular:

i) We questioned the Regge formulas used in ACGL and CGL, which did not respect factorization. Thus, we proposed a Regge parametrization, that, according to Caprini, Colangelo, Gasser and Leutwyler [8] (CCGL) violated crossing symmetry rather strongly. Later on, we “re-discovered” the existing data on $\pi\pi$ total cross sections: they turned out to be well described with our Regge formulas, but not with those of ACGL and CGL. Furthermore, and although the issue is now irrelevant because there is data to compare with, we have shown that crossing is satisfied with our Regge equations.

We have also shown that the D2 wave parameterization in ACGL, CGL, derived in the seventies from intermediate energy data, was used outside its applicability range.

ii) Since the CGL phase shifts did not satisfy a number of sum rules when using a D2 wave and a Regge description compatible with data, we proposed a “tentative solution” [3], from a fit to an average of data. Surprisingly, CCGL claimed that this “tentative solution” did not describe experiment. We have shown here that this only happens because ACGL, CGL and CCGL only consider a subset of all published data.

iii) We have shown here that the uncertainty imposed by ACGL and CGL on the difference between P and S0 phase shifts at 800 MeV, is largely underestimated when taking into account all published data and their systematic uncertainties, that are emphasized in some of the original references. This is an important input for their Roy equation analysis and, as remarked by CCGL, the main source of error for the Olsson sum rule.

iv) CCGL claim that the CGL Roy equation analysis remains unchanged (except for the S2 wave) if using our Regge asymptotics. We agree that this can be the case for certain observables, like the S and P wave scattering lengths (which are indeed compatible with our values). However, other quantities, like the phase shifts at intermediate energy or the D-wave scattering lengths can vary considerably.

v) We have confirmed our result for the b_1 P-wave threshold parameter with a new sum rule that depends little on Regge asymptotics, or the precise form of the P wave parametrization.

vi) Finally, we have recently shown that the CGL S and P phase shifts fail to satisfy three forward dispersion relations up to $\sqrt{s} \leq 800$ MeV by several standard deviations.

Apart from this discussion, we also used forward dispersion relations to check the consistency [5] of different data sets, including the precise data from kaon decays. As it is well known, there are many different phase shift analyses, often incompatible even within the same experiment. Surprisingly, we found that some of the most frequently used data sets are inconsistent

with forward dispersion relations and sum rules, and should therefore be used with caution. Finally, these forward dispersion relations and sum rules were used to constrain the data fits and to provide a consistent amplitude easy to implement by those interested in $\pi\pi$ scattering.

Acknowledgments

Work partially supported by DGICYT Spain, contracts FPA2005-02327 and BFM2002-01868, and the EURIDICE network contract HPRN-CT-2002-00311 as well as the EU Hadron Physics Project, contract number RII3-CT-2004-506078.

REFERENCES

1. Ananthanarayan, B., Colangelo, G., Gasser, J., and Leutwyler, H., *Phys. Rep.*, **353**, 207, (2001).
2. Colangelo, G., Gasser, J., and Leutwyler, H., *Nucl. Phys.* **B603**, 125, (2001).
3. Peláez, J. R., and Ynduráin, F. J., *Phys. Rev.* **D68**, 074005 (2003).
4. Peláez, J. R., and Ynduráin, F. J., *Phys. Rev.* **D69**, 114001 (2004).
5. Peláez, J. R., and Ynduráin, F. J., *Phys. Rev.* **D71**, 074016 (2005).
6. Peláez, J. R., and Ynduráin, F. J., hep-ph/0411334. AIP Conf. Proc.756:48-59,2005.
7. de Trocóniz, J. F., and Ynduráin, F. J., *Phys. Rev.*, **D65**, 093001, (2002), *Phys. Rev.* **D71**, 073008 (2005).
8. Caprini, I., Colangelo, G., Gasser, J., and Leutwyler, H., *Phys. Rev.* **D68**, 074006 (2003).
9. Ynduráin, F. J., hep-ph/0310206. eConf30614:4,2003.
10. Colangelo, G., hep-ph/0501107.
11. Biswas, N. N., et al., *Phys. Rev. Lett.*, **18**, 273 (1967) Cohen, D. et al., *Phys. Rev.* **D7**, 661 (1973). Robertson, W. J., Walker, W. D., and Davis, J. L., *Phys. Rev.* **D7**, 2554 (1973). Hanlon, J., et al, *Phys. Rev. Lett.*, **37**, 967 (1976); Abramowicz, H., et al. *Nucl. Phys.*, **B166**, 62 (1980)
12. Martin, B. R., Morgan, D., and and Shaw, G. *Pion-Pion Interactions in Particle Physics*, Academic Press, New York (1976).
13. Losty, M. J., et al. *Nucl. Phys.*, **B69**, 185 (1974); Hoogland, W., et al. *Nucl. Phys.*, **B126**, 109 (1977); Cohen, D. et al., *Phys. Rev.* **D7**, 661 (1973); Durusoy, N. B., et al., *Phys. Lett.* **B45**, 517 (1973).
14. Hyams, B., et al., *Nucl. Phys.* **B64**, 134, (1973).
15. Grayer, G., et al., *Nucl. Phys.* **B75**, 189, (1974).
16. Hyams, B., et al., *Nucl. Phys.* **B100**, 205 (1975).
17. Au, K. L., Morgan, D., and Pennington, M. R. *Phys. Rev.* **D35**, 1633 (1987).
18. Estabrooks, P., and Martin, A. D., *Nucl. Physics*, **B79**, 301, (1974).
19. Kamiński, R., Lesniak, L, and Rybicki, K., *Z. Phys.* **C74**, 79 (1997) and *Eur. Phys. J.* **C4**, 4 (2002).
20. Protopopescu, S. D., et al., *Phys. Rev.* **D7**, 1279, (1973).

- 21. R. Kamiński, L. Leśniak and B. Loiseau, *Phys. Lett.* **B551** .
- 22. Donoghue, J. F., Gasser, J., and Leutwyler, H., *Nucl. Phys.*, **B343**, 341 (1990); Ynduráin, F. J., *Phys. Lett.* **B578**, 99 (2004) and (E), **B586**, 439 (2004); Ananthanarayan, B., et al., *Phys. Lett.* **B602**, 218 (2004); Ynduráin, F. J., *Phys. Lett.* **B612**, 245-257 (2005).